

# CALCIUM SC PLUS

Reg. No. B3659 Act No. 36 of 1947



FERTILISER

## FERTILISER GROUP 2

**A calcium suspension concentrate for foliar application with added zinc and boron.**

Calcium	(Ca)	242 g/kg	(400 g/l)
Boron	(B)	2 g/kg	( 3 g/l)
Zinc	(Zn)	18 g/kg	( 30 g/l)

**SG = 1,65 ± 0.02 @ 20°C**

Content • Batch number • Manufacture date



**Registered, Manufactured and Distributed by:**  
Ag-Chem Africa (Pty) Ltd (Reg. No. 1998/013411/07)  
P.O.Box 589, Silverton, 0127, Gauteng, RSA.  
Tel: (012) 803-0145 • Fax: (012) 803-8418  
[www.agchem.co.za](http://www.agchem.co.za)

CALCIUM SC PLUS: 01-2013/12

**SHAKE WELL** Shake well before use • Store at room temperature  
Keep out of direct sunlight • Keep container closed when not in use



LIFT HERE

- **CALCIUM SC PLUS** should preferably be applied early in the early morning or late afternoon.
- It is advisable to do a miscibility test prior to mixing **CALCIUM SC PLUS** with other chemicals.
- Consult with your chemical distributor should crop specific programmes or any other information be required regarding the use of **CALCIUM SC PLUS**.

#### DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	MAXIMUM RATE PER HECTARE	RATE PER 100 ℓ WATER	REMARKS
Apples	5 ℓ	200 – 250 mℓ	Apply from petal fall to one month before harvest with 10 – 14 day intervals. Apply as a full cover spray in 1 000 – 3 000 ℓ water/ha. Avoid run-off.
Avocados and Mangoes	5 ℓ	250 mℓ	Apply from fruit set to one month before harvest, with 10 – 14 day intervals. Apply in 2 000 ℓ water/ha.
Bananas	2 ℓ	-	Apply every 10 – 14 days to one month before harvest. Apply in 30 ℓ water/ha.
Citrus	5 ℓ	125 mℓ	Apply from fruit set. Do not apply within one month before harvest. Apply in 4 000 ℓ water/ha.
Cucurbits (Cucumbers, Melons)	2 – 4 ℓ	-	Apply with 7 – 10 day intervals from first fruit set to one month before harvest. Repeat if necessary. Apply in 200 – 400 ℓ water/ha.
Deciduous Fruits	5 ℓ	250 mℓ	Apply with 7 – 14 day intervals from petal fall to one month before harvest. Apply in 2 000 ℓ water/ha. If conditions are conducive to fruit russet (variety, weather conditions, etc) start applications 6 weeks after petal fall.
Grapes	5 ℓ	-	Apply every 10 – 14 days from fruit set in 1 000 ℓ water/ha.
Groundnuts	3 – 5 ℓ	-	Start application at 4 – 6 leaf stage. If necessary, repeat application with 10 – 14 day intervals. Apply in 30 – 200 ℓ water/ha.
Nuts (Macadamias, Pecans)	5 ℓ	250 mℓ	Start application 7 days after petal fall. Repeat if necessary with 10 – 14 day intervals. Do not apply within one month before harvest. Apply in 1 000 – 2 000 ℓ water/ha.
Peppers (field grown)	5 ℓ	-	Apply from 4 – 6 leaf stage to one month before harvest. Repeat applications every 7 – 10 days. Apply in 50 – 500 ℓ water/ha.
Strawberry (field grown)	3 – 5 ℓ	-	Start application early in the new season until one month before harvest. Apply 3 applications with 10 – 14 day intervals in 300 – 500 ℓ water/ha.
Tomato (field grown)	5 ℓ	-	Start application at the end of flowering on second truss until one month before harvest. If necessary repeat with 7 – 10 day intervals. Apply a maximum of 5 applications per season. Apply in 50 – 500 ℓ water/ha.
Vegetables (field grown) (Cabbage, Carrots, Lettuce, etc.)	3 – 5 ℓ	-	Apply from 4 – 9 leaf stage until one month before harvest. Repeat with 10 – 14 day intervals. Apply in 50 – 500 ℓ water/ha.