

COPSTAR

Reg. No. B3487 Act No. 36 of 1947



FERTILISER

FERTILISER GROUP 2

COPSTAR is a liquid suspension concentrate fertiliser for use as a foliar feed to maintain or correct copper levels in plants.

Copper (Cu) 102 g/kg (120 g/ℓ)

SG = 1,20 ± 0.02 @ 20°C

Content • Batch number • Manufacture date



Registered, Manufactured and Distributed by:
Ag-Chem Africa (Pty) Ltd (Reg. No. 1998/013411/07)
P.O.Box 589, Silverton, 0127, Gauteng, RSA.
Tel: (012) 803-0145 • Fax: (012) 803-8418
www.agchem.co.za

COPSTAR: 01-2013/12

SHAKE WELL Shake well before use • Store at room temperature
Keep out of direct sunlight • Keep container closed when not in use



LIFT HERE

For optimum results **COPSTAR** should be used in a programme and in combination with other balanced fertilizers and foliar feeds.

COPSTAR is compatible with most agricultural remedies. It is **however advisable to do a miscibility test prior to mixing with other chemicals.**

COPSTAR should preferably be applied in the early morning or late afternoon. **Do not apply to plants that are undergoing a period of moisture or heat stress.**

Consult with your chemical distributor should crop specific programmes or any other information be required about the use of **COPSTAR.**

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	MAXIMUM RATE PER HECTARE	RATE PER 100 ℓ WATER	REMARKS
Grapes	5 ℓ	350 ℓ	Apply with bud swell and repeat application at 75 % petal fall. Apply in a minimum of 1 000 – 3 000 ℓ water per hectare.
Citrus	10 ℓ	350 ml	Apply in September, before flowering. Repeat 3 – 4 weeks later if necessary. Apply as a high volume application in 1 000 – 3 000 ℓ water per hectare.
Potatoes, Tomatoes, Beans and Cruciferae	5 ℓ	350 ml	Apply 1 – 2 applications early in the season, when there is enough leaves to absorb the spray. Repeat every 7 – 10 days if necessary. Depending on plant size apply in 500 – 1 500 ℓ water per hectare.
Onions	3 ℓ	600 ml	Apply from ghoif ball size bulbs, and repeat 14 days later. Apply in a minimum of 500 ℓ water per hectare.
Other crops	3 ℓ	350 ml	Apply every 10 – 14 days if necessary to correct copper deficiencies.

Use lower rates as a maintenance spray and higher rates to correct deficiencies.