

EASYCROP

Reg. No. K6483 Act No. 36 of 1947



FERTILISER GROUP 1

A water soluble 5:10:2 (26) fertiliser with magnesium, sulfur and micro-elements for foliar feeding and fertigation.


Nitrogen (N)	78 g/kg	*Iron (Fe)	3510 mg/kg
Phosphorus (P)	149 g/kg	*Manganese (Mn)	702 mg/kg
Potassium (K)	31 g/kg	Molybdenum (Mo)	78 mg/kg
Sulphur (S)	41 g/kg	*Zinc (Zn)	450 mg/kg
Magnesium (Mg)	31 g/kg	*EDTA chelated	
Boron (B)	1027 mg/kg		
*Copper (Cu)	56 mg/kg		



agchem  Member of **Rolfes**
GROUP

Registered, Manufactured and Distributed by:
Ag-Chem Africa (Pty) Ltd (Reg. No. 1998/013411/07)
P.O.Box 589, Silverton, 0127, Gauteng, RSA.
Tel: (012) 803-0145 • Fax: (012) 803-8418
www.agchem.co.za

Content • Batch number • Manufacture date



www.agchem.co.za

EASYCROP-02-2014/03



EASY CROP is a completely water-soluble fertilizer and chlorine free.

EASY CROP is readily taken up by the plant and is formulated for use as both a foliar feed and for fertigation.

EASY CROP should preferably be applied in the early morning or late afternoon. **Do not apply to plants that are undergoing a period of moisture or heat stress.**

Consult with your chemical distributor, consultant or manufacturer should crop specific programs or any other information be required around the use of.

EASY CROP is compatible with most agricultural remedies. **It is however advisable to do a miscibility test prior to mixing with other chemicals.**

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	APPLICATION RATE	REMARKS
Citrus	500 g/100 ℓ water	Apply during periods of stress.
Dry beans	4 kg/ha	Start application at the 8 leaf stage. Repeat every 14 days for a total of 3 applications.
Maize	4 kg/ha	Start application 4 – 6 weeks after emergence.
Mangoes	500 g/100 ℓ water	Apply 2 weeks after bud break and repeat 14 days later.
Potatoes	4 kg/ha	Apply after 80% emergence. Repeat every 14 days for a total of 5 applications.
Soya beans	4 kg/ha	First application at 8 leaf stage and repeat 14 days later.
Sugarcane	4 kg/ha	Apply 12 – 16 weeks after harvest.
Sunflowers	5 kg/ha	Apply 4 – 6 weeks after emergence and repeat 14 days later.
Tobacco	4 kg/ha	Start application 4 weeks after transplant. Repeat every 7 days for a total of 3 applications.
Vegetables	4 kg/ha	Apply after 80% emergence. Repeat every 14 days for a total of 5 applications.
Wheat	4 kg/ha	Apply 6 – 8 weeks after emergence.

The rate of application is dependent on the soil fertility and reaction required. The lower rates should be used for maintenance while the higher rates will correct nutritional deficiencies.