





FERTILISER GROUP 2

Reg. No. K5242 Act No. 36 of 1947

A water soluble 3:1:6 (46) fertiliser with micro-elements for foliar feeding and fertigation.



Nitrogen	(N)	141 g/kg
Phosphorus	(P)	46 g/kg
	$(P_{2}O_{5})$	105 g/kg
Potassium	(K)	275 g/kg
	$(K_{2}0)$	331 g/kg
Sulphur	(S)	13 g/kg
Magnesium	(Mg)	2 g/kg

Zinc*	(Zn)	500	mg/kg
Iron*	(Fe)	500	mg/kg
Manganese*	(Mn)	200	mg/kg
Copper*	(Cu)	200	mg/kg
Boron	(B)	1000	mg/kg
Molybdenum	(Mo)	50	mg/kg
Cytokinins		34	mg/kg
Auxins		57	mg/kg

Content • Batch number • Manufacture date

*FDTA chelated





Registered, Manufactured and Distributed by: Ag-Chem Africa (Pty) Ltd (Reg. No. 1998/013411/07) P.O.Box 589, Silverton, 0127, Gauteng, RSA. Tel: (012) 803-0145 • Fax: (012) 803-8418 www.agchem.co.za



EASYGRO Flower & Fruit is recommended for use during plant growth periods that require relatively high potassium levels such as during flowering, fruit formation and fruit growth.

EASYGRO Flower & Fruit is a completely water-soluble fertilizer and chlorine free.

EASYGRO Flower & Fruit is readily taken up by the plant and is formulated for use as both a foliar feed and for fertigation.

For optimum results EASYGRO Flower & Fruit should be used in a program in combination with the other EASYGRO range of fertilizers.

EASYGRO Calmag + TE* should be used in combination with EASYGRO Flower & Fruit for increased fruit quality, shelf life and for correcting calcium and magnesium deficiencies.

EASYGRO Flower & Fruit is compatible with most agricultural remedies. It is however advisable to do a miscibility test prior to mixing with other chemicals

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	RATE PER HECTARE	RATE PER 100 LITRE Water	REMARKS
Foliar feed	1 – 3 kg	1 – 2 kg	Vegetable and ornamental crops. Apply as a full cover spray. Apply at $1-2$ week intervals throughout the flowering and fruiting period.
	1-3 kg	1 – 3 kg/30 ℓ water	$\mbox{\bf Aerial Application.}$ Apply at $1-2$ week intervals throughout the flowering and fruiting period.
	4 – 6 kg	1 – 2 kg	
Fertigation	5 – 10 kg	0,25 – 2 kg	Use lower concentrations for seedlings and increase the rate as the plants mature. Use during periods that require a high nitrogen level.

The rate of application is dependent on the soil fertility and reaction required. The lower rates should be used for maintenance while the higher rates will correct nutritional deficiencies.

EASYGRO Flower & Fruit should preferably be applied in the early morning or late afternoon. Do not apply to plants that are undergoing a period of moisture or heat stress.

Consult with your chemical distributor, consultant or manufacturer should crop specific programs or any other information be required around the use of the **EASYGRO** range of fertilisers.

*Easygro Calmag + TE Reg. No. K5243, Act No. 36 of 1947 Ag-Chem Africa (Pty) Ltd.