

# SIZE-UP SC

Reg. No. K9656 Act No. 36 of 1947



## FERTILISER GROUP 1

**A water soluble liquid suspension fertilizer for foliar feeding and fertigation.**

Phosphorus	(P)	139 g/kg	(226 g/ℓ)
Potassium	(K)	176 g/kg	(287 g/ℓ)

**SG (@ 20 °C) = 1.63 ± 0.02**

Content • Batch number • Manufacture date



**Registered, Manufactured and Distributed by:**  
Ag-Chem Africa (Pty) Ltd (Reg. No. 1998/013411/07)  
P.O.Box 589, Silverton, 0127, Gauteng, RSA.  
Tel: (012) 803-0145 • Fax: (012) 803-8418  
[www.agchem.co.za](http://www.agchem.co.za)

[www.agchem.co.za](http://www.agchem.co.za)



**PRECAUTIONS:**

- **SIZE-UP SC** should preferably be applied in the early morning or late afternoon. **Do not apply to plants that are undergoing a period of moisture or heat stress.**
- Wear suitable protective clothing.
- Clean all equipment before and after use.
- Destroy empty container and do not use it for any other purpose.
- Keep away from children, uninformed people, animals and foodstuffs.
- Wash hands after use.

**DIRECTIONS FOR USE: Use only as directed**

- Do not exceed the appropriate application rates.
- To be used only when recommended by a suitably qualified person.
- For optimum results, **SIZE-UP SC** should be used in a programme and in combination with other balanced fertilisers and foliar feeds.
- Consult with your chemical distributor, consultant or the registration holder should crop specific programmes or any other information be required.

**MIXING INSTRUCTIONS:**

Always follow the label recommendations. Shake container thoroughly before use. Any dosing equipment must be carefully calibrated to ensure correct and even application.

Partially fill tank with water, begin agitation. Add required amount of **SIZE-UP SC**. Add other chemicals if compatible. Fill tank and agitate thoroughly.

**COMPATIBILITY:**

Read all labels carefully. It is advisable to do a miscibility test prior to mixing with other chemicals. Check with your supplier before tank mixing.

**STORAGE:**

Keep in a cool dry chemical store, out of reach of children and livestock. Store in a well-sealed container away from sunlight. Protect from cold and heat extremes.

CROP	MAXIMUM RATE PER HECTARE	RATE PER 100 ℓ WATER	REMARKS
<b>Bean Crops</b> (Soybean, Groundnut and Green Beans)	2 ℓ	500 mℓ	Apply as a full cover spray.
<b>Cabbage crops, Carrots</b>	2 ℓ	500 mℓ	Apply as a full cover spray.
<b>Citrus</b>	15 ℓ	200 – 500 mℓ	Apply as a full cover spray.  <b>Fruit colour and size:</b> Apply the first application after petal fall. Apply 2 – 3 applications per season. Do not apply within 30 days before harvest. Use the lower dosage on soft citrus. Application should take place in early morning when plants are not undergoing heat stress. Apply another 1 – 2 applications postharvest.
<b>Cereals</b> (Wheat, Barley, Rice, Rye, Oats)	2 ℓ	500 mℓ	Apply as a full cover spray.
<b>Cotton</b>	2 ℓ	500 mℓ	Apply as a full cover spray.

CROP	MAXIMUM RATE PER HECTARE	RATE PER 100 ℓ WATER	REMARKS
Cucurbits (Pumpkins, Melons, Cucumber etc.)	2 ℓ	500 mℓ	Apply as a full cover spray.
Deciduous Fruit (Apples, Pears, Apricots, Peaches, Cherries, Fig, Nectarine, Plums, Pomegranates)	15 ℓ	200 – 500 mℓ	Apply 2 – 3 applications after petal fall. <b>Fruit colour and size:</b> Apply the first application four weeks prior to harvest and apply a second application 2 weeks later. Apply another 1 – 2 applications postharvest.
Wine Grapes	15 ℓ	400 – 750 mℓ	Apply 1 – 2 applications prior to flowering; typically at variation and 7 days later. Apply another 1 – 2 applications postharvest.
Table Grapes	15 ℓ	400 – 750 mℓ	Apply 1 – 2 applications prior to flowering; typically at variation and 7 days later. <b>Berry colour and size:</b> Apply the first application four weeks prior to harvest and apply a second application 2 weeks later. Apply another 1 – 2 applications postharvest.
Flowers and Ornamentals	2 ℓ	200 mℓ	Apply as a full cover spray.
Olives	10 ℓ	500 mℓ	Apply as a full cover spray.
Lucerne and Pasture	2 ℓ	500 mℓ	Apply as a full cover spray.
Macadamias and Nut Crops	2 ℓ	500 mℓ	Apply as a full cover spray.
Maize	2 ℓ	500 mℓ	Apply as a full cover spray.
Potatoes	2 ℓ	500 mℓ	Apply as a full cover spray.
Onions	2 ℓ	500 mℓ	Apply as a full cover spray.
Subtropical Fruit (Avocados, Litchi, Mangoes, Coffee, Papaya)	2 ℓ	500 mℓ	Apply as a full cover spray. <b>Fruit colour and size:</b> Apply the first application after petal fall. Apply 2 – 3 applications per season. Do not apply within 30 days before harvest. Application should take place in early morning when plants are not undergoing heat stress. Apply another 1 – 2 applications postharvest.
Sugarcane	2 ℓ	500 mℓ	Apply as a full cover spray.
Tomatoes, Peppers	5 – 10 ℓ	500 mℓ	Apply during the reproductive stages of the plants. The dosage rate depends on the cultivar and adequate nutritional levels during the specific growth stage.
Turf	2 ℓ	500 mℓ	Apply as a full cover spray.
Other crops not mentioned	2 ℓ	500 mℓ	Apply as a full cover spray.