

SKINSET

Reg. No. B5110 Act No. 36 of 1947

FERTILISER GROUP 2

SKINSET is a liquid suspension concentrate fertiliser containing elemental sulphur and calcium. It is for use as a soil application to maintain or correct sulphur and calcium levels in soils and plants.

Calcium	(Ca)	62 g/kg	(90.5 g/ℓ)
Sulphur	(S)	317 g/kg	(463 g/ℓ)

SG @ 20°C = 1,46 ± 0,02

Content • Batch number • Manufacture date

agchem  Member of **Rolfes**
GROUP

Registered, Manufactured and Distributed by:

Ag-Chem Africa (Pty) Ltd (Reg. No. 1998/013411/07)

P.O.Box 589, Silverton, 0127, Gauteng, RSA.

Tel: (012) 803-0145 • Fax: (012) 803-8418

www.agchem.co.za

SKINSET: 07-2016



SHAKE WELL Shake well before use • Store at room temperature
Keep out of direct sunlight • Keep container closed when not in use



LIFT HERE

RECOMMENDATIONS FOR USE:

For optimum results **SKINSET** should be used in a programme and in combination with other balanced fertilisers and foliar feeds.

It is advisable to do a miscibility test prior to mixing with other chemicals.

Consult with your chemical distributor should crop specific programmes or any other information be required about the use of **SKINSET**.

Ensure thorough agitation of the product during filling and spraying. Always use spray tank strainers when pouring product into spray tank. The spray tank must be flushed out with clean water at the end of the day.

Ensure good agitation throughout application.

Use lower rates as maintenance and higher rates to correct deficiencies.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	MAXIMUM RATE PER HECTARE	REMARKS
Citrus	50 – 120 ℓ	Apply through the irrigation system. Will correct calcium and sulphur levels in root zone. Will also replace excessive sodium in root zone.
Legumes (Alfalfa, Soybeans, Groundnuts)	50 – 100 ℓ	Apply through the irrigation system as a pre-plant soil treatment. Shallow incorporation in soil is important for efficacy of product
Maize	4 – 6 ℓ	Apply as an in-furrow application during planting.
	50 – 100 ℓ	Apply to soil before planting and incorporate.
Onions	50 – 100 ℓ	Apply to soil before planting and incorporate.
Other Crops	50 – 100 ℓ	Apply through the irrigation system. Will correct calcium and sulphur levels in root zone. (Contact an Ag-Chem Africa representative for recommendations).
Potatoes	100 ℓ	Apply in a minimum of 300 ℓ water/ha as a direct spray during planting in-furrow. Can be applied directly onto seed tubers. Do not apply on seed tubers with a high incidence of pests and diseases which affect the quality of the seed tubers.
Vines	50 – 100 ℓ	Apply through the irrigation system as a pre-plant soil treatment.
Wheat	4 – 6 ℓ	Apply as an in-furrow application during planting
	50 – 100 ℓ	Apply to soil before planting and incorporate.